



Rev Mex Med Forense, 2017, 3(2):91-98

ISSN: 2448-8011

Thiel Soft-Fix method for long term preservation

Review article

Edmundo Denis-Rodríguez ¹, Ángel Augusto Aguirre-Gutiérrez ²

SUMMARY

Long term cadaveric preservation represents a medical challenge, considering the toxicological consequences of formaldehyde in the environment and personal health status; stiffness of the tissues increases the difficulty of their manipulation in medical school and teaching hospitals. In recent years a new method of long-term preservation has emerged; it was proposed by Walter Thiel, known as Thiel Soft-fix method; it uses low concentrations of formaldehyde, also avoiding tissue stiffness for longer periods. In this review, we analyze the features of Thiel Soft-Fix method for long-term preservation of cadaver and tissues.

Keywords: Embalming, Preservation, Thiel Soft-Fix method

Received: May 15th, 2018, Accepted: June 5th, 2018, Online: August 15th, 2018

¹ Master in Forensic Medicine, Doctor of Forensic Sciences, Institute of Forensic Medicine, Universidad Veracruzana

² Master in Forensic Medicine, Doctor of Forensic Sciences, Institute of Forensic Medicine, Universidad Veracruzana

Corresponding author: Edmundo Denis-Rodríguez, eddenis@uv.mx

INTRODUCTION

The long term preservation of corpses represents a medical challenge. Traditional embalming uses high concentrations of preservatives, especially formaldehyde (Okada, 2012). From numerous studies, formaldehyde has been related to various pathologies, both acute (skin irritation and conjunctiva, respiratory distress) and chronic (chronic dermatitis, corneal opacity, chronic respiratory and renal alterations). In addition to the aforementioned pathologies, special mention deserves the relationship between the prolonged use of formaldehyde and the appearance of malignant neoplasms, especially hematological ones (Hayashi, 2016).

In addition to the individual and environmental toxicological implications, it is known that the traditional technique of long-term preservation hardens the bodily tissues in an important way, which is why it is difficult to manipulate them by medical students in Faculties of Medicine and Teaching Hospitals (Hammer, 2015). From the above, the importance of having a long-term preservation method in which the tissues retain most of their pre-mortem consistency without reducing the preservative efficacy becomes important.

The Thiel Soft-Fix method is an embalming technique originally proposed by the Austrian anatomist Walter Thiel (Anatomy Institute of Graz, Austria) (Thiel, 1992). After several years of improvement, the original publication was

made in a German-language magazine, which is why it was not generally known at first. 10 years later a second article was published in English, which has led to the implementation of this technique in various centers around the world (Thiel 2002).

The technique of Walter Thiel, known as Thiel Soft-Fix Method, has been gaining popularity in the world (Hammer, 2015). In 2011, a study was conducted in which they observed this technique is known in 53% of the departments of anatomy, dissection and forensic sciences in the world, although its routine use as an embalming method is only done in 10% of the included centers (Benkhadra, 2011b). Its main use is in Europe, although in recent years it has begun to be used in other countries, such as India, Thailand, Japan and Argentina (Sangchay, 2014).

SCIENTIFIC BASIS

Modern embalming uses several preservatives, but the most frequent is formaldehyde; although surfactants and modifiers are currently used to improve the cosmetic result of this technique, when corpses require embalming for prolonged periods, high concentrations of formaldehyde are required (Benkhadra 2011). As a result, tissues dehydrate, harden and change their color and general appearance, making it difficult to use in Anatomy, Surgery and Interventionist Techniques. Figure 1 shows the tissue preserved in a traditional way with formaldehyde.



Figure 1. Tissue preserved with the traditional formaldehyde technique at high concentrations.

The Thiel Soft Fix technique uses preservatives other than formaldehyde (in fact, although it contains it, its concentration is very low) (Thiel, 2002). It is a technique designed so that tissues, especially the muscular and integumentary types, retain most of their premortem texture and tone (Hassan 2014). In a study carried out on muscle and connective tissue, it was observed that in the samples taken from embalmed cadavers with the

Thiel Soft Fix technique, the muscle tissue presented focal degradation and loss of fiber distribution, unlike non-embalmed corpses or embalmed corpses. with formaldehyde (Benkhadra, 2011); This observation suggests that muscular and connective changes could explain the soft consistency observed in the Thiel Soft Fix technique (Joy, 2015, Liao, 2015). Figure 2 shows the tissue preserved by the Thiel method.



Figure 2. Tissue preserved using the Thiel Soft-Fix method.

TECHNIQUE DESCRIPTION

The success of the Thiel Soft-Fix technique depends on the complete and orderly completion of the stages that

compose it (Eisma, 2013b). This technique has two stages (Bertone, 2011):

Initial Embalming

After washing and draining the entire vascular tree, to eliminate clots and any other factor that decreases the circulation of the fluid, the injection solution is introduced by the cervical route (internal carotid or common carotid artery), which is obtained by combining the Solution A and solution B (proposed by Thiel) and adding 300 ml of formaldehyde and 700 g of sodium sulfite, for a total volume of 15.7 liters. Solution A contains 3% boric acid, 30% ethylene glycol, 20% ammonium nitrate, 5% potassium nitrate and 42% water for a usual total volume of 14-15 liters. Solution B contains 10% ethylene glycol and 1% 4-chloro-3-methylphenol for a total volume of 500 ml. This same mixture is introduced to the body orally and rectally to try to incorporate it into the digestive tract and respiratory tree.

Immersion

Once the injection solution was incorporated into the cadaver, it is placed in a pool or tub, which contains the so-called Immersion Solution. This solution contains 10% of ethylene glycol, 2% of formaldehyde, 2% of solution B of Thiel, 3% of boric acid, 10% of ammonium nitrate, 5% of potassium nitrate, 7% of sodium sulfite and 65% % of water, for a total volume that will depend on the size of the pool, but enough to submerge the body in its entirety. The recommended total immersion time is 30 days, which allows it to retain its characteristics even after repeated use (Munirama, 2016). Once past this period, the body is removed from the pool and placed in a bag with airtight seal; if signs of dehydration are observed over time, the body can be submerged again in the same basin for a period of 5-7 days (Thiel, 2002). Even body fragments can be introduced into the immersion solution and after a period of 2-3 months they recover

their original tonicity in vivo (Hunter, 2014).

CURRENT USE

Cadavers preserved using the Thiel Soft-Fix method have been used in various areas of the interventional medical field. In general, its soft consistency has allowed Medicine and Surgery students to perform dissection practices in conditions very similar to those they would present in a living corpse. In fact, studies have been carried out in which the subjective experience of medical students is evaluated using the preserved cadaver with the Thiel Soft-Fix method (Balta, 2015).

Cadavers preserved with this technique have been used in the practice and assessment of interventional diagnostic studies, such as urinary tract endoscopy (Bele, 2016) and magnetic resonance (Guerguieva, 2014). It has also been used in the practice of surgical or invasive procedures such as renal transplantation (Cabello, 2015), abdominal laparoscopy (Eisma, 2011), thyroid surgery (Eisma, 2013a), urological procedures (Healy, 2015), surgery oral (Hölzle, 2012), vascular microsurgery (Obedescu, 2014), nephrectomy (Prasad, 2012; Rai, 2015) and orotracheal intubation (Szücs, 2016). Finally, they have been used in various studies of tissue physiology, such as biomechanical studies of the tendinous tissue (Fessel, 2011), studies of the acoustic mechanics of the middle ear (Guignard, 2011) and determination of the mechanical properties of cortical bone (Unger, 2010). It is evident that with the increase of the worldwide knowledge about this technique, the diagnostic and surgical applications will be more and more numerous.

CONCLUSION

For more than 60 years, the traditional technique of embalming uses several preservatives; the most frequently used is formaldehyde; this substance has been linked to various pathologies, both acute and chronic, which give it its obviously toxic nature.

When it is intended to embalm a corpse for a long time, whether for academic or didactic purposes, high concentrations of formaldehyde are required, which increases its risk of toxicity to the embalmer.

Various techniques have arisen in which minimal or no formaldehyde quantities are used for long-term preservation. Since 1992, a technique called Thiel Soft-Fix has been proposed; prolonged preservation of a corpse is achieved without the toxic effects of formaldehyde and, as an additional benefit, conserving the texture of the tissues in vivo almost completely.

In Mexico this type of technique has not been implemented; consequently, the response that the corpses can have, histologically, to a humid warm medium has not been evaluated either. For more than 20 years the use of the Thiel Soft-Fix technique has been implemented worldwide, which is a technique that allows the long-term preservation of corpses, with a minimum and non-toxic amount of formaldehyde and with the benefit additional to allow tissues to retain their soft consistency similar to in vivo conditions.

It is convenient the implementation in our environment of techniques like the one mentioned, with no toxicity for the embalmer. Also, once it is implemented, the histological changes that occur in the embalmed corpse should be studied, both to know objectively the changes that are made in a humid warm climate and to understand histologically the reason why this technique ensures tissues of the body to maintain a similar appearance to their premortem condition.

REFERENCES

1. Balta JY, Lamb C, Soames RW (2015). A pilot study comparing the use of Thiel- and formalin-embalmed cadáver in the teaching of human anatomy. *Anat Sci Educ*, 8(1): 86-91.
2. Bele U, Kelc R (2016). Upper and lower urinary tract endoscopy training on Thiel-embalmed cadavers. *Urology*, 93: 27-32
3. Benkhadra M, Bouchot A, Gerard J (2011). Flexibility of Thiel's embalmed cadavers: the explanation is probably in the muscles. *Surg Radiol Anat*, 33(4): 365-8.
4. Benkhadra M, Gerard J, Genelot D (2011b). Is Thiel's embalming method widely known? A world survey about its use. *Surg Radiol Anat*, 33(4): 359-63.
5. Bertone VH, Blasi E, Ottone NE (2011). Walther Thiel Method for the preservation of corpses with maintenance of the main physical properties in vivo. *Rev Arg Anat*, 2(3): 71-100
6. Cabello R, Gonzalez C, Quicios C (2015). An experimental model for training in renal transplantation surgery with human cadavers

- preserved using W. Thiel's embalming technique. *J Surg Educ*, 72(2): 192-7.
7. Eisma R, Gueorguieva M, Immel E (2013). Liver displacement during ventilation in Thiel embalmed human cadavers - a possible model for research and training in minimally invasive therapies. *Minim Invasive Ther Allied Technol*, 22(5): 291-6.
8. Eisma R, Lamb C, Soames RW (2013b). From formalin to Thiel embalming. What Changes? One anatomy department's experiences. *Clin Anat*, 26(5): 564-71.
9. Eisma R, Mahendran S, Majumdar S (2011). A comparison of Thiel and formalin embalmed cadavers for thyroid surgery training. *Surgeon*, 9(3): 142-6.
10. Fessel G, Frey K, Schweizer A (2011). Suitability of Thiel embalmed tendons for biomechanical investigation. *Ann Anat*, 193(3): 237-41.
11. Gueorguieva MJ, Yeo DT, Eisma R (2014). MRI of Thiel-embalmed human cadavers. *J Magn Reson Imaging*, 39(3): 576-83.
12. Guignard J, Stieger C, Kompis M (2013). Bone conduction in Thiel-embalmed cadaver heads. *Hear Res*, 306: 115-22.
13. Hammer N, Löffler S, Bechmann I (2015). Comparison of modified Thiel embalming and ethanol-glycerin fixation in an anatomy environment: Potentials and limitations of two complementary techniques. *Anat Sci Educ*, 8(1): 74-85.
14. Hassan S, Eisma R, Harry LE (2014). Surgical training of anastomotic technique using Thiel cadaver. *J Plast Reconstr Aesthet Surg*, 67(10): 250-1.
15. Hayashi S, Naito M, Kawata S (2016). History and future of human cadaver preservation for surgical training. *Anat Sci Int*, 91(7): 1-7.
16. Healy SE, Rai BP, Biyani CS (2015). Thiel embalming method for cadaver preservation: a review of new training model for urologic skills training. *Urology*, 85(3): 499-504.
17. Hölzle F, Franz EP, Lehm Brock J (2012). Thiel embalming technique: a valuable method for teaching oral surgery and implantology. *Clin Implant Dent Relat Res*, 14(1): 121-6.
18. Hunter A, Eisma R, Lamb C (2014). Thiel embalming fluid: a new way to revive formalin-fixed cadaveric specimens. *Clin Anat*, 27(6): 853-5.
19. Joy J, Mcleod G, Lee N (2015). Quantitative assessment of Thiel soft-embalmed human cadavers using shear wave elastography. *Ann Anat*, 202: 52-6.
20. Liao X, Kemp S, Corner G (2015). Elastic properties of Thiel-embalmed human ankle tendon and ligament. *Clin Anat*, 28(7): 917-24.
21. Munirama S, Eisma R, Columb M (2016). Physical properties and functional alignment of soft-embalmed Thiel human cadáver when used as a simulator for ultrasound-guided regional anaesthesia. *Br J Anaesth*, 116(5): 699-707.
22. Obedescu A, Moubayed SP, Harris PG (2014). A new microsurgical model using Thiel-embalmed arteries and comparison of two suture techniques. *J Plast Reconstr Aesthet Surg*, 67(3): 389-95.

23. Okada R, Tsunoda A, Momiyama N (2012). Thiel's method of embalming and its usefulness in surgical assessments. *Nihon JGK*, 115(8): 791-4.
24. Prasad B, Tang B, Eisma R (2012). A qualitative assessment of human cadavers embalmed by Thiel's method used in laparoscopic training for renal resection. *Anat Sci Educ*, 5(3): 182-6.
25. Rai BP, Stolzerburg JU, Healy S (2015). Preliminary validation of Thiel embalmed cadavers for laparoscopic radical nephrectomy. *J Endourol*, 29(5): 595-603.
26. Sangchay N (2014). The soft cadaver (Thiel's method): The new type of cadaver of Department of Anatomy, Siriraj Hospital. *Siriraj Med H*, 66: 228-231.
27. Szücs Z, Lázló CJ, Baksa G, (2016). Suitability of a preserved human cadaver model for the simulation of facemask ventilation, direct laryngoscopy and tracheal intubation: a laboratory investigation. *Br J Anaesth*, 116(3): 417-22.
28. Thiel W (1992). The preservation of the whole corpse with natural color. *Ann Anat*, 174(3): 185-95.
29. Thiel W (2002). Supplement to the conservation of an entire cadaver according to W. Thiel. *Ann Anat*, 184(3): 267-9.
30. Unger S, Blauth M, Schmoelz W (2010). Effects of three different preservation methods on the mechanical properties of human and bovine cortical bone. *Bone*, 47(6): 1048-53.
31. Verstraete MA, Van Der Straeten C, De Lepeleere B (2015). Impact of drying and thiel embalming on mechanical properties of achilles tendons. *Clin Anat*, 28(8): 994-1001.
32. Wilke HJ, Werner K, Häussler K (2011). Thiel-fixation preserves the non-linear deformation characteristic of spinal motion segments, but increases their flexibility. *J Mech Behav Biomed Mater*, 4(8): 2133-7.

